

BROILERS

AGE	VACCINE	ROUTE
Day 1 HATCHERY	NCD + IB	Spray
Day 9	IBD (Gumboro)	Drinking Water
Day 12	NCD + IB	Eye Drop/Drinking Water
Day 18	IBD (Gumboro)	Drinking Water

LAYERS

AGE	VACCINE	ROUTE
Day 1 HATCHERY	1. Mereks 2. NCD + IB 3. IBD (Gumboro)	Subcutaneous Spray Beak Dip
Day 7	NCD + IB	Eye Drop/Drinking Water
Day 12	IBD (Intermediate)	Drinking Water
Day 21	IBD (Intermediate)	Drinking Water
Day 28	NCD (Lasota)	Eye Drop/Drinking Water
6 Weeks	Fowl Pox	Wing Web
8 Weeks	NCD + IB	Eye Drop/Drinking Water
9 Weeks	Deworming	Drinking Water
10 Weeks	Fowl Typhoid	Intra-Muscular
12 Weeks	Debeaking (If Necessary)	
15 Weeks	Deworming	Drinking Water
16 Weeks	Fowl Typhoid	Intra-Muscular
18 Weeks	NCD + IB Inactivated	Drinking Water

NOTE:

- Repeat NCD (Lasota) after 38-40 weeks, then every 6-8 weeks.
- Repeat deworming every 4-6 weeks.
- Always transport vaccines in ice.
- Give multivitamins before and after vaccination.
- Vaccination should be in a clean environment by a trained person.
- Do not use water which contains any disinfectant or soap.

FOR ORDERS: Call  or WhatsApp  +256709852604